

Pansenstimulans Bolus

Rumen Stimulant Bolus

agilan® Complementary feed

Mineral feed

For cattle

Composition

Dicalcium phosphate, monosodium phosphate, yeast, potato starch, skimmed milk powder



Analytical constituents

Crude protein	12.7 %
Crude fiber	1.7 %
Crude fat	3.7 %
Crude ash	45.4 %
Calcium	9.8 %
Sodium	2.3 %
Phosphorus	8.7 %
Magnesium	1.2 %

Content of additives per kg

Vitamins:

Vitamin B ₁ (3a821)	800 mg
Vitamin B ₂ (3a825i)	2,500 mg
Vitamin B ₆ (3a831)	2,000 mg
Vitamin B ₁₂ / Cyanocobalamin	16,000 µg
Vitamin K ₃ (MNB) (3a711)	1,600 mg
Niacin (3a314)	10,000 mg
Calcium-D-pantothenate (3a841)	6,000 mg
Folic acid (3a316)	400 mg
Biotin (3a880)	40,000 µg

Microorganisms:

Saccharomyces cerevisiae CNCM I-1077

(Levucell SC) (E1711)

100 x 10⁹ cfu

Feeding recommendation

1-2 pieces (each 95 g) per animal per day, administer orally with commercial donors.

Feeding instructions

The amount of *Saccharomyces cerevisiae* in the daily ration must not exceed 8.4 x 10⁹ cfu at 100 kg body weight in dairy cows or 4.6 x 10⁹ cfu in beef cattle. For each additional 100 kg of body weight, 1.8 x 10⁹ cfu should be added to dairy cows and 2 x 10⁹ cfu for beef cattle.

Pansenstimulans Bolus

Rumen Stimulant Bolus

Recommended use

Pansenstimulans Bolus (Rumen Stimulant Bolus) with live yeast as well as high levels of vitamins of the entire B-complex and selected phosphorus carriers. Pansenstimulans Bolus (Rumen Stimulant Bolus) optimizes the activity of the microorganisms in the rumen and thus the use of the nutrients offered. Since all water-soluble vitamins can be stored only in small quantities by the cow, a continuous sufficient supply via the own synthesis is only possible with a non-contaminated metabolic and optimal rumen function.

Storage

Store in a dry place. Seal container well after use, so that there is no moisture. Shelf life guarantee is only valid for unopened packages at dry, cool storage (2 - 25 °C).

Date of revision of the text: July 2021